

# Seed Notes

## for Western Australia

No. 9 Acacia

### IN THIS ISSUE

This issue of **Seed Notes** will cover the genus *Acacia*.

- Description
- Geographic distribution and habitat
- Reproductive biology
- Seed collection
- Seed quality assessment
- Seed germination
- Recommended reading

## Acacia

The name *Acacia* comes from the Greek *acacia*, *ace* or *acis* meaning a point or thorn, or from *acazo*, to sharpen, although this name applies more to African than Australian species (Australian *Acacia* have no thorns or larger prickles, unlike those that are native to Africa). Many species of *Acacia*, or wattles as they are commonly called in Australia, are valuable for a range of uses, in particular as garden plants. They are also used for amenity plantings, windbreaks, shade trees, groundcovers, erosion and salinity control. The timber of some *Acacia* is very hard and is ideal for fence posts (e.g. *A. saligna* or jam). Other *Acacia* species are used to make furniture and ornaments. The seed of some wattles is a good food source for birds, other animals and humans as 'bush tucker'. *Acacia* are generally fast growing, but many are short-lived.



*Phyllodes and flowers of Acacia aprica.*

*Photo – Andrew Crawford*

### Description

In Australia, *Acacia* (family Mimosaceae) are woody plants that range from prostrate under-shrubs to tall trees. *Acacia* flowers are small, regular and usually bisexual. They occur in spikes or in globular heads and range in colour from cream to intense yellow. The leaves of *Acacia* may be bipinnate (the primary leaflets being again divided into secondary leaflets) or entirely absent at maturity and the petiole

is modified to form a leaf-like structure or phyllode. These phyllodes may be flat or terete. Some species do not have phyllodes and the flattened stems or cladodes act as leaves. Foliage can vary from blueish to dark green to silvery grey. Most species of *Acacia* have glands on the axis of the phyllodes, although in Australian species these appear to have no function, but aid in the identification of the species.



Department of Environment and Conservation



Our environment, our future



Top: *Acacia trulliformis*. Above: *Acacia alata* var. *platyptera*. Above right: *Acacia insolita* ssp. *recurva*. Photos – Anne Cochrane

## Geographic distribution and habitat

*Acacia* is the largest genus of higher plants in Australia with more than 1200 species worldwide and more than 900 of these occur in Australia over a wide range of habitats. *Acacia* occur in a range of climatic and soil conditions, from coastal situations to mountain tops, from tropical to arid inland areas. They grow in a range of soil types from limestone to laterite, from granite to deep sands. They are generally drought hardy. The genus is not thought to be susceptible to the dieback disease, *Phytophthora cinnamomi*, although many species are threatened by rising salt levels and waterlogging. The centre of diversity for *Acacia* in Western Australia is in the heavily cleared agricultural wheatbelt, where in addition to salinisation, small population size and weed invasion further threaten the survival of these species.



Approximate distribution of *Acacia* in Australia.





*Acacia nervosa*.

Photo – Andrew Crawford

## Reproductive biology

Many species of *Acacia* flower at an early age, and are insect pollinated, although wind may have a lesser role to play in reproduction. Birds may also play a part in pollination, attracted to the flowers by the insects and by extra-floral nectaries on the phyllodes. There may be some pollinator specificity. Although there are usually large numbers of flowers per inflorescence, few pods are formed. The fruit is a legume that may be straight, curved or coiled. Multiple seeds are contained within the fruits. These seeds contain a protein-rich appendage, or aril, on the outside of the



seed which may be large or small and can range in colour from greenish to white to yellow and to red. The arils attract birds and ants, and are known to disperse the seed of many species. This myrmecochory, or ant distribution of seed, is well established in Australia and for some species, burial of seed beneath the soil surface prevents destruction of seed during wildfires. Seedlings can often be seen germinating from old ants' nests after fire. Birds are also known to disperse seed. Natural hybridisation between species occurs to some extent, but is not widespread like in the eucalypts. Seed of the genus are known to have a long storage life and some species are reported to still be viable after 50 or more years when stored under ambient temperatures. Drying and freezing of seed will further improve this longevity.

Clockwise from far left: Buds of *Acacia*; multiple seedlings of *Acacia* emerging after fire; *Acacia microneura* flowers; and *Acacia insolita* ssp. *recurva* inflorescence.

Photos – Anne Cochrane

## Seed collection

Collection of *Acacia* seed is very easy and seed is generally readily available. The seed is ripe when the pod dries out, changes colour and becomes brittle and splits. Seed of the majority of *Acacia* species in Western Australia ripen in late spring to early summer after mid year flowering, but ripening will occur quicker in more northern hotter climates (e.g. north of Perth) compared to those species from the southern part of the State (e.g. Albany). The pods can split quite quickly so collection needs to be well timed to coincide with the shedding. Time from bud formation to fruit ripening is often a year, with plants displaying small buds at the same time as seed shed. On maturity, the seed is hard and ranges from light brown to glossy black depending on the species. It is possible to collect seed of some species of *Acacia* when fully formed but still green for immature sowing. However, for longevity of seed it is preferable to collect only mature seed.



Top: *Acacia dorsenna*.

Photo – Anne Cochrane

Above: Collecting an *Acacia* from Dandaragan.

Photo – Tony Friend



## Seed quality assessment

Acacia seeds are often predated by insects. These insects lay their eggs in the developing fruit. The larvae grow as the fruit develops and the seeds are damaged. Birds are also found to attack the fruits, and grasshoppers, in particular locusts, will damage young green fruits. Any damaged seed should be removed and a simple float test can be conducted to distinguish good from bad seed. The damaged or light seed will float and the whole seed will sink. After placing it in water make sure that you dry the seed off very well before it is stored.



Clockwise from top left: Open pod of *Acacia* species; furry pods of *Acacia alata* var. *platyptera*, *Acacia insolita* ssp. *efoliolata*; *Acacia alata* var. *platyptera* fruits opening to release seed; *Acacia cyclops* with red arils; immature fruits of *Acacia congesta* ssp. *wonganensis*.

Photos – Anne Cochrane



Both insect damaged and undamaged seed of *Acacia pharangites*.



*Acacia aprica* seed.



*Acacia clydonophora* seed.



5mm

*Acacia ataxiphylla* ssp. *magna* seed.



Left: *Acacia leptalea* seed and pod.

Below: Insect damage on *Acacia insolita* ssp. *efoliolata* pods.

Photos – Anne Cochrane



## Germination of seeds

Most species are hard-seeded and require pre-treatment to the seed coat before germination will occur. Heat shock (wet or dry) or manual scarification of the seed coat (chipping of a portion of the hard coat to reveal the endosperm, abrading with an emery board or file, or placing the seed in a paint or jewellery tumbler with small stones or sand) are all methods used to artificially reduce the hardness of seeds. Chipping the seed coat is the most successful method, but time consuming, and the use of a microscope may



be required if the seed is small. The immersion of seed in hot to boiling water reduces the hardness of seeds in many species, although there are some species that have softer seed coats and very hot water can kill these seeds. These latter species will generally germinate without treatment. Be aware that there is a wide variation in the heating requirement for different species. There are many similarities between treatment of *Acacia* seeds and those of other hard-seeded species such as *Daviesia*, *Gastrolobium*, *Gompholobium* and *Chorizema*.



Above: Germinating seed of *Acacia*.

Above left: Seed of *Acacia drummondii* ssp. *elegans*.

Far left: *Acacia drummondii* ssp. *elegans* seed in the fruit.

Left: *Acacia insolita* seed in the fruit.

Photos – Anne Cochrane



## Seed Notes

for Western Australia



These **Seed Notes** aim to provide information on seed identification, collection, biology and germination for a wide range of seed types for Western Australian native species.



They have been written and compiled by Anne Cochrane, Manager of DEC's Threatened Flora Seed Centre.

Concept by Grazyna Paczkowska.

Designed by DEC's Graphic Design Section.

The **Seed Notes** are available from [www.naturebase.net](http://www.naturebase.net)

## Seed Notes

are published by the Perth Branch of the Wildflower Society of Western Australia (Inc.) with assistance from the Western Australian Lotteries Commission and the Department of Environment and Conservation (DEC).



The rare *Acacia aprica*. Photo – Sally Madden



*Acacia teretifolia*. Photo – Andrew Crawford



*Acacia glaucoptera* fruit. Photo – Anne Cochrane

## Recommended reading

Allen, O. N. and Allen, E. K. 1981. *The Leguminosae*. A source book of characteristics, uses and nodulation. University of Wisconsin Press, Wisconsin.

Cavanagh, H. T. 1987. Germination of hard-seeded species (Order Fabales). In *Germination of Australian Native Plant Seed*. P. J. Langkamp (ed) Pp. 58-70. Inkata Press, Melbourne.

Elliot, W. R. and Jones, D. L. 1984. *Encyclopaedia of Australian Plants Suitable for Cultivation*. Volume 2. Lothian Publishing, Melbourne.

*Flora of Australia* 11A and 11B. Mimosaceae. Acacia. ABR/CSIRO, Melbourne.

Sharr, F. A. 1978. *Western Australian Plant Names and their Meanings. A Glossary*. University of Western Australia Press, Perth.

Simmons, M. 1981. *Acacias of Australia*. Nelson, Melbourne.